

## Germination Studies in *Rhanterium epapposum* Oliv

Majda Khalil Suleiman, Narayana Ramachandra Bhat,  
Mehdi Saleh Abdal, Sameeha Zaman, Sheena Jacob and Rini Rachel Thomas

Arid land Agriculture and Greenery Department,  
Kuwait Institute for Scientific Research, P.O. Box 24885, 13109 Safat, Kuwait

**Abstract:** Conservation of native plants are possible by their utilization in urban landscape, as indigenous plant species have evolved and adapted to the local harsh climatic conditions over the years and they are more likely to function adequately under the local climate as opposed to exotic plants. As Kuwait's native plants are threatened in their natural habitat and have begun disappearing at an alarming rate, their use in landscape projects will help in conserving biodiversity and heritage. Dormancy is widespread among desert shrubs, with only a few species having non-dormant seeds. Effective dormancy breaking treatments are important to develop mass propagation techniques in native plants like *Rhanterium epapposum*. In this germination study, effects of dry heat (50°C) and/or various concentrations (500, 750 and 1000 ppm) of gibberellic acid (GA<sub>3</sub>) on seed germination of *Rhanterium epapposum* were assessed. Treating the seeds with 750 ppm of GA<sub>3</sub> was found to be effective for *Rhanterium epapposum* with 71% of germination. The next best treatment combination (52%) was exposure to dry heat (50°C) for ten days followed by a treatment with 500 ppm GA<sub>3</sub> for 24 hrs.

**Key words:** Urban landscape % Greenery development % Dormancy % Germination study % Conservation of native plants

### INTRODUCTION

Kuwait has a typical desert climate with extreme temperatures, minimal and variable rainfall, intense sunshine and frequently occurring dust storms. The climate is characterized by extremely hot summers, with daytime temperature exceeding 50°C and winter, cooler and at times wet, with temperature sometimes falling below 4°C [1]. The rainy season extends from October to May. The mean annual rainfall is 113mm [2]. The total conventional freshwater resources available in Kuwait are six million cubic meter per year, while the total water demand has exceeded 350 million m<sup>3</sup>/yr in 2000 [3]. Native soil in Kuwait is mostly sandy in texture with high infiltration rate and is calcareous in nature [3].

Perennial vegetation cover is sparse, usually less than 10% [4]. Although introduced ornamental plants are necessary in urban and sub-urban landscape for diversity, they rely on high levels of nutrients and water. Conservation of native plants are possible by their utilization in urban landscape, as indigenous plant species have evolved and adapted to the local harsh climatic conditions over the years and they are more likely to

function adequately under the local climate as opposed to exotic plants. As Kuwait's native plants are threatened in their natural habitat and have begun disappearing at an alarming rate, their use in landscape projects will help in conserving biodiversity and heritage [5]. Shrubs, in particular, are significant and have good potential for urban landscape application [6]. Unfortunately, efficient propagation and establishment techniques that are crucial for both conservation of native plants and large-scale use in landscape programs are currently unavailable.

Accordingly, 'arfaj' (*Rhanterium epapposum* Oliv, Family: Asteraceae) was selected for its aesthetic appeal, flower color and potential to adapt to urban landscape conditions. It is the national flower of Kuwait. It is a C<sub>3</sub> desert shrub that can form monotonous stands covering vast areas of north-eastern Arabia [4]. It grows as a perennial woody shrub approximately 80 cm high with many stems branching out from the base. The leaves are small and narrow and in late spring, it is covered with straw-yellow flowers about 1cm wide. It flowers from April to May and produces numerous fruits (Fig. 1) which forms in late spring and falls off the branches after maturity. It accumulates under the shrub and remains dormant until




Fig. 1: *Rhanterium epapposum* (a) capituli; (b) seeds

favorable conditions for germination prevail. Each fruit (capitulum) contains about 6-8 seeds that are transported by wind or water. The total area covered by this plant is about 2% of Kuwait. It is considered one of the main desert forage plants for camels and sheep [4].

Germination is a critical phase in the life cycle of a plant, particularly in arid regions where seeds can be confronted by severe environmental constraints such as high temperatures, drought and high levels of salinity as a result of fluctuating soil moisture [4]. Baskin and Baskin [7] Stated that dormancy is widespread among desert shrubs, with only a few species having non-dormant seeds. In Kuwait and adjacent areas, *Rhanterium epapposum* is found mainly in deep sandy soils [8]. The capitulum is the unit of dispersal for *Rhanterium epapposum*, either being blown short-distances by the wind or transported over long distances by zoochory [4]. For germination to take place, the capituli must be positioned with their basal part on the ground, in an upright position [9]. He also demonstrated that dormancy over a number of years is a typical feature of the seeds. Often, several seeds germinate at once in a capitulum and it appears that mortality of seedlings due to intra-specific competition may not be the norm [9]. The regeneration of *Rhanterium epapposum* from seed has not been observed, although this must be the natural method of propagation [8]. Seed germination under simulated environmental conditions ranged from 0-4.69% [10]. It is difficult to propagate and establish this species by vegetative means. It can be propagated by tissue culture method [11] but, the production cost will be higher than the seedling method. In view of the above facts, studies were conducted to determine the effects of pre-treatments on germination of *Rhanterium epapposum* seeds.

## MATERIALS AND METHODS

**Seeds:** Seeds used in this study were obtained from seed bank of Kuwait Institute for Scientific Research (KISR) in November 2007. These seeds were collected from Kuwait desert in 2004. Prior to their use in the experiment, seed viability was determined using Triphenyl Tetrazolium Chloride (TTC) Test [12]. In this test, viable embryos are stained pink due to the reduction of 2, 3, 5- TTC by reparative activity in the cell. For this, 100 seeds (four replicates of 25 seeds each) were soaked in distilled water overnight, excised to expose the embryos, placed in a petri dish, soaked in 0.1% TTC solution, covered the petri dish with aluminum foil and kept for 24 hours at  $25 \pm 1^\circ\text{C}$ . Following the treatment, seeds were washed thoroughly with distilled water to remove excess stain and examined under the microscope.

**Germination Studies:** Seeds stored at room temperature were kept in an oven at  $50^\circ\text{C}$  for 10 days or 20 days and treated with various concentrations of gibberellic acid ( $\text{GA}_3$ ) for 24 hours. Treatments included exposure to dry heat at  $50^\circ\text{C}$  for 10 or 20 days and/ or pre-sowing soaking in  $\text{GA}_3$  solution (0, 500, 750 or 1,000 ppm) for 24 hours. There were twelve treatments, which were replicated five times in a completely randomized design. Each replication contained 20 seeds. The control seeds were not subjected to either heat or  $\text{GA}_3$  treatment. The total germination was calculated when no more seeds germinated. Pre-treated seeds were sown in agricultural soil medium in Petri dishes and maintained in growth chamber at  $18^\circ\text{C}$  with 60% humidity.

The data were analyzed using R analysis procedure of Analysis of Variance (ANOVA) and Duncan's Multiple Range Test to ascertain the significant differences among treatments [13, 14].

## RESULTS AND DISCUSSION

Viability of seeds used in these studies was found to be 88%. The highest germination of (Table 1) 71% was obtained from seeds that were treated with 750 ppm  $\text{GA}_3$  for 24 hours (Fig. 2). The next best treatment combination (52%) was the ten days of dry heat ( $50^\circ\text{C}$ ) exposure followed by a treatment with 500 ppm  $\text{GA}_3$  for 24 hrs. When the seeds were subjected to 10 days heat treatment, increase in the concentration of  $\text{GA}_3$  had a negative effect on the germination. Exposure of seeds to dry heat alone

Table 1: Effect of Dry Heat (50°C) Exposure and GA<sub>3</sub> on Germination of *Rhanterium epapposum* Seeds

Heat Treatment	Germination (%) <sup>i</sup>			
	GA <sub>3</sub> Treatment (ppm)			
	0	500	750	1000
RT <sup>ii</sup>	50±10.75 de <sup>iii</sup>	21±3.28 abc	71±5.36 e	44±2.60 cd
50°C 10 Days	27±5.39 abcd	52±5.39 de	38±3.89 bcd	27±6.71 abcd
50°C 20 Days	7±2.28 a	17±9.11 ab	28±10.99 abc	22±9.93 abc

<sup>i</sup> Seeds with 2 mm or longer radicle or shoot are considered as germinated.

<sup>ii</sup> RT- Room Temperature (25°C)

<sup>iii</sup> The means followed by the same letter are not statistically different at p#0.001




Fig. 2: *Rhanterium epapposum* germination

for 10 or 20 days also did not improve the germination. Untreated seeds resulted in 50% of germination. Extended duration (20 days) of dry heat exposure diminished the germination. The improvement in germination by pre-treatment was significant at P#0.001 level.

The germination percentage of 71% in the present study is a clear improvement over the previous reports where only 4.69% [10] germination was recorded under simulated environmental conditions. The fact that, pre-treatment with GA<sub>3</sub> increased the germination percentage suggests that the dormancy in this species may be due to physiological factors. GA<sub>3</sub> treatment alters hormonal balance in favor of promoters to trigger germination process and shorten time needed for germination [15]. Results of this study are important as desert rehabilitation and biodiversity conservation is crucial to prevent the extinction of valuable native plants in Kuwait [2, 5].

## CONCLUSIONS

Effective dormancy breaking treatments are important to develop mass propagation techniques in native plants like *Rhanterium epapposum* which exhibit dormancy to adapt to the harsh environmental conditions in the desert. Pre-treating the seeds with 750 ppm GA<sub>3</sub> was effective in breaking seeds dormancy and in significantly improving the germination in *Rhanterium epapposum*.

## ACKNOWLEDGEMENT

The authors would like to thank the Kuwait Foundation for the Advancement of Sciences (KFAS) and Kuwait Institute for Scientific Research for providing funds and encouragement during the investigation.

## REFERENCES

1. FAO., 2004. AQUASTAT- FAO's Information System on Water and Agriculture. <http://www.fao.org/ag/agl/aglw/aquastat/countries/kuwait>. 15/11/06.
2. Omar, S.A.S., Y. Al-Mutawa and S. Zaman, 2007. Vegetation of Kuwait, Kuwait: Kuwait Institute for Scientific Research, pp: 17-39.
3. Hamoda, M.F., 2001. Desalination and water resource management in Kuwait. Desalination, 138(1-3):165.
4. Brown, G., 2001. Vegetation ecology and biodiversity of degraded desert areas in north-eastern Arabia-Habilitation Thesis. Rostock University, pp: 90-92.
5. Suleiman, M.K., N.R. Bhat and R. Bellen, 2004. Various attempts in greening the State of Kuwait. Arch. Agron. Soil Sci., 50: 353-360.

6. AboEl-Nil, M., L. Al-Sabah, H. Al-Menaie, S. Bastaki and S. Zaman. 1993. Mass propagation of stress tolerant desert plants by tissue culture: Phase I. Kuwait Institute for Scientific Research Report No. KISR 4300, Kuwait.
7. Baskin, C.C. and J.M. Baskin, 1998. Seeds. Ecology, biogeography and evolution of dormancy and germination. Academic Press, San Diego.
8. Halwagy, R., A.F. Moustafa and S.M. Kamel, 1982. On the ecology of the desert vegetation in Kuwait. *J. Arid Environ.*, 5: 95-107.
9. Thalen, D.C., 1979. Ecology and utilization of desert shrub rangelands in Iraq. Junk, The Hague.
10. Taha, F.K., S.A. Omar, A.E. Hegab, S. Zaman and A.A. Nasser, 1988. Range management project Phase II. Assessment of range lands in Kuwait and improvement for forage animal production and de-desertification. Kuwait Institute for Scientific Research, Report No. KISR 2762, Kuwait.
11. Sudharsan, C., M. AboEl-Nil and J. Hussain 2003. Tissue culture technology for conservation and propagation of certain native plants of Kuwait. *J. of Arid Environ.*, 54: 133-147.
12. ISTA (International Seed Testing Association), 1999. International rules for seed testing. *Seed Sci. Technol.*, 27: 333.
13. Little T.M. and F.J. Hills, 1978, *Agricultural Experimentation Design and Analysis*. (U.S.A: John Wiley and Sons Inc.), pp: 63-65.
14. Crawley, M.J., 2005. *Statistics- An Introduction using R*. England: John Wiley Sons Ltd., pp: 155-185.
15. Ergenoglu, F., S. Tangolar, S. Gok and A.B. Kuden, 1997. The effects of some pretreatments for promoting germination of dormant grape seeds. *Acta. Hortic.*, 441: 201-212.